

HOSTED BY



ELSEVIER

Available online at www.sciencedirect.com

ScienceDirect

www.elsevier.com/locate/bsci

Review article

Interactions between sleep, stress, and metabolism: From physiological to pathological conditions



Camila Hirotsu*, Sergio Tufik, Monica Levy Andersen

Department of Psychobiology, Universidade Federal de São Paulo, São Paulo, Brazil

ARTICLE INFO

Article history:

Received 19 November 2014

Received in revised form

15 September 2015

Accepted 16 September 2015

Available online 28 September 2015

Keywords:

Sleep

Stress

Metabolism

Cortisol

Hypothalamic–pituitary–adrenal axis

Obesity

ABSTRACT

Poor sleep quality due to sleep disorders and sleep loss is highly prevalent in the modern society. Underlying mechanisms show that stress is involved in the relationship between sleep and metabolism through hypothalamic–pituitary–adrenal (HPA) axis activation. Sleep deprivation and sleep disorders are associated with maladaptive changes in the HPA axis, leading to neuroendocrine dysregulation. Excess of glucocorticoids increase glucose and insulin and decrease adiponectin levels. Thus, this review provides overall view of the relationship between sleep, stress, and metabolism from basic physiology to pathological conditions, highlighting effective treatments for metabolic disturbances.

© 2015 Brazilian Association of Sleep. Production and Hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Sleep and stress interact in a bidirectional fashion, sharing multiple pathways that affect the central nervous system (CNS) and metabolism, and may constitute underlying mechanisms responsible in part for the increasing prevalence of metabolic disorders such as obesity and diabetes [1]. Hormones like melatonin and others from the hypothalamic–pituitary–adrenal (HPA) axis modulate the sleep–wake cycle, while its dysfunction can disrupt sleep. In turn, sleep loss influence the HPA axis, leading to hyperactivation [2]. In the first part of this paper, we focus on the definitions of sleep and the HPA axis, and the relationship between sleep and

stress. In the second part, we review the effects of sleep and stress on the metabolism, addressing mainly sleep deprivation, circadian alterations, and key sleep and stress disorders. Finally, we connected these topics to provide a better understanding of the intrinsic relationship between sleep, stress and metabolism, and suggest possible targets for future intervention.

The secretory activity of the HPA axis follows a distinct 24 h pattern. CRH is released in a circadian-dependent and pulsatile manner from the parvocellular cells of the PVN [3]. In fact, the circadian rhythm of cortisol secretion derives from the connection between the PVN and the central pacemaker, the suprachiasmatic nucleus (SCN) [4]. The close

*Correspondence to: Rua Napoleao de Barros, 925, Vila Clementino - SP 04024-002, São Paulo, Brazil. Tel.: +55 11 5908 7193; fax: +55 11 5572 5092.

E-mail address: milahirotso@gmail.com (C. Hirotsu).

<http://dx.doi.org/10.1016/j.slsci.2015.09.002>

1984-0063/© 2015 Brazilian Association of Sleep. Production and Hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Fig. 1 – 24-h individual cortisol profile showing the minimum (nadir), the maximum (acrophase), the onset of the circadian rise, and the amplitude of the cortisol profile. After a nadir during the early night, there is an important rise in ACTH and cortisol in the late night, reaching a peak near the awakening time, driven by circadian oscillators, such as sleep.

proximity of AVP-containing SCN nerve endings near CRH-containing neurons in the PVN suggests that via this projection circadian information is imprinted onto the HPA-axis [5]. Typically, the nadir (time point with the lowest concentration) for cortisol occurs near midnight. Then, cortisol levels increase 2–3 h after sleep onset, and keep rising into the waking hours. The peak happens in the morning at about 9 a. m. [4]. Along the day, there is a progressive decline that is potentiated by sleep, until it reaches the nadir and the quiescent period (Fig. 1). In general, 3 main pathways are essential for biological clock function: the input (*zeitgebers*, retina) → SCN circadian pacemaker (as clock genes, neurotransmitters, peptides) → output (pineal melatonin synthesis, thermoregulation, hormones). Then, these factors interact with the sleep–wake cycle to modulate, for example, sleep propensity and sleep architecture, and influence behavior, performance or hormonal output such as cortisol [4].

2. Disturbed or shifted sleep, sleep loss and HPA axis

Many stressful situations, such as low socioeconomic status and chronic work overload, have been associated with a deficit in sleep duration and several neuroendocrine effects (for review, see [6]). Indeed, there is long-standing evidence of reciprocal interactions between the HPA axis and sleep regulation [7], which will be discussed below.

Circadian misalignment affects sleep architecture and may also reduce total sleep time. Both advanced and delayed phases result in disruption of the normal phase relationship between SWS and REM sleep [8]. During the first day of an 8 h phase delay, profound disruptions in the 24 h cortisol rhythm were found, with a higher nadir value mediated by the lack of the inhibitory effects caused by sleep onset, and lower acrophase values due to the lack of the stimulatory effects of awakening, resulting in an overall 40% reduction in the

rhythm [9]. Five days after the shift, the cortisol profile had adapted to the new schedule [9]. On the other hand, an advanced phase of 8 h had advanced the timing of the cortisol nadir by about 3 h and 20 min, with marked reduction in the quiescent period, and increased the rising phase of cortisol secretion by 3 h [10]. In this last case, no adaptation of the timing of the acrophase to the new schedule was observed. In summary, these studies confirm that the misalignment of the sleep–wake cycle has a negative impact on the stress system. Although it seems to be a short-term effect probably due to a biphasic pattern of the cortisol rise after the shift, it may also contribute to metabolic changes. Alterations of the HPA axis may play a causative role in sleep disorders such as insomnia. HPA axis dysfunction may be secondary to a clinical sleep disorder, such as obstructive sleep apnea (OSA), leading to other complications.

Insomnia is a sleep disorder characterized by difficulties in falling or staying asleep or having restorative sleep, associated with daytime impairment or distress [11]. Despite the relationship between sleep and the HPA axis, little is known about the neurobiological basis of this sleep disorder and its link with HPA axis activation. One study did not show any significant differences in urinary cortisol between control and poor sleepers [12]. However, another study presented a positive correlation between polysomnographic indices of sleep disturbance and urinary free cortisol in adults with insomnia [13]. Patients with insomnia without depression do present high levels of cortisol, mainly in the evening and at sleep onset, suggesting that, rather than the primary cause of insomnia, the increase in cortisol may be a marker of CRH and norepinephrine activity during the night [14]. Preceding evening cortisol levels are correlated with the number of the following night's nocturnal awakenings, independent of insomnia [15]. However, excessive activation of the HPA axis induces sleep fragmentation [16], while the sleep fragmentation increases cortisol levels [15], suggesting that the HPA axis may contribute to the initiation as well as the perpetuation of chronic insomnia [15]. There is still debate whether the activation of the HPA axis found in insomnia is secondary to sleep loss or a marker of CRH activity.

OSA is a common sleep disordered breathing, characterized by recurrent apneas (complete breathing cessation) or hypopneas (shallow breathing), upper airway constriction, hypoxemia, hypercapnia, autonomic activation, and EEG arousal and sleep fragmentation, leading to daytime fatigue and sleepiness [17]. As nocturnal awakening is associated with pulsatile cortisol release and autonomic activation, we can expect OSA to lead to HPA axis activation through the same mechanisms involved in arousal and sleep fragmentation [4]. However, the studies to date are contradictory. Some have shown that continuous positive airway pressure (CPAP) therapy for OSA does not lower cortisol while the acute withdrawal of CPAP does not change cortisol levels [18]. On the other hand, other authors have demonstrated that CPAP does reverse hypercortisolemia [19]. A systematic review revealed that only 2 studies showed statistically significant differences in cortisol levels after CPAP treatment [20].

Elevated cortisol levels were reported in patients with OSA by some studies [21], but not in others [22]. Responsiveness of ACTH to CRH administration was much higher in obese

patients with OSA, possibly due to alterations in the central control of ACTH secretion and impairment in the negative feedback of glucocorticoids [23]. A recent study showed that serum basal and peak cortisol levels were lower in OSA patients when compared to the control group during 1 μ g ACTH and glucagon stimulation tests, showing an association between OSA and hypocortisolemia in the morning with reduced responses to ACTH and glucagon stimulation tests [24]. Many of the discrepancies observed in the literature are reflective of methodological differences. The majority of studies are limited by assessment of cortisol at a single time point. The available studies do not provide clear evidence regarding whether OSA is associated with alterations in cortisol levels or that treatment with CPAP changes cortisol levels. Methodological concerns such as infrequent sampling, failure to match comparison groups on demographic factors known to impact cortisol levels (age, body mass index etc.), and inconsistent control of confounding factors may have limited the findings. However, there is evidence that excessive HPA axis activation may be a result from sleep loss, hypoxemia, and autonomic activation, playing an important role in the metabolic alterations arising from OSA [17].

Many studies have shown increase in cortisol levels during the nighttime period of total sleep deprivation and in the prolonged wakefulness of the following day. This is likely a result of stress due to the effort of maintaining wakefulness, as high frequency EEG activity is correlated with indices of arousal and cortisol release [25,26]. However, some authors also reported no change [27,28] or a decrease in cortisol levels [29,30] after 1 or more nights of sleep deprivation. These discrepancies seem to be influenced by insufficient frequency of blood sampling, small sample size, and by fatigue and sleepiness. In animals, however, the results are more consistent. Adult rats subject to paradoxical sleep deprivation during 96 h show increased levels of corticosterone, which are normalized after 48 h of sleep rebound [31]. Notwithstanding, it is important to consider that animal models do not accurately reflect human physiology; and thus, it is difficult to compare these results.

Studies using chronic protocols of sleep restriction, which model a widespread condition in modern society, have also addressed the role of HPA axis. The first study assessed the effect of 6 consecutive nights of 4 h in bed in young men, showing increased levels of cortisol in the afternoon and early evening, and a shorter quiescent period, with onset delayed by 1.5 h [16]. The rate of decrease of free cortisol in saliva was nearly 6 times slower in sleep restricted volunteers compared to fully rested condition. Notably, chronic short sleepers do present higher levels of cortisol compared to chronic long sleepers [32].

Sleep deprivation seems to be related to the elevation of cortisol, reflecting impairment of HPA axis regulation, and resulting in glucocorticoid overload, which can lead to large deleterious effects on the body. Moreover, there is an association between short sleep duration and higher risk of developing obesity and type II diabetes, suggesting the HPA axis hyperactivation as one of the mechanism involved in the metabolic consequences of sleep loss [33,34].

3. Effects of glucocorticoids on sleep and metabolism

Classic studies in rats and humans have demonstrated that exogenous CRH is able to modulate sleep by increasing EEG frequency and wakefulness and decreasing SWS [35,36]. However, studies focused on the direct effects of glucocorticoids have shown that they increase time spent awake at the expense of REM sleep [37]. Other studies show that cortisol decreases SWS when MRs are activated, while dexamethasone increases awakening after activation of GRs [38]. The effects of both exogenous and endogenous glucocorticoids on sleep EEG depend on the type and location of the receptors activated (MR vs. GR), the dose of cortisol/corticosterone used, and the optimal cortisol levels to effect maximal nocturnal CRH suppression [4,39]. Studies that have demonstrated decreases in SWS with elevated cortisol levels and total occupation of GR may be due to excessive GR activation in the amygdala. The effects are opposite to the known inhibitory action found in PVN and anterior pituitary, leading to positive feedback [39,40]. However, the effects of glucocorticoids as well as CRH on REM are not well understood and most of them are contradictory [4].

Chronic exposure to excess glucocorticoids, such as occurs during diseases like Cushing's syndrome, can offer insight into the role of these hormones in sleep. For example, consistent alterations in polysomnographic recordings are reported in Cushing's syndrome, such as reduction of SWS, increased sleep latency, enhanced wake time, shortened REM latency, and elevated REM density, among others [17], reflecting the deleterious effects of glucocorticoid excess. Bierwolf and colleagues [41] have demonstrated that adrenal secretory activity starts predominantly during periods of NREM in both Cushing's and healthy patients, showing a link between pituitary-adrenal activity and the ultradian rhythmicity of NREM and REM sleep.

Reductions in sleep duration have become common due to the socioeconomic demands and opportunities in modern society [42]. In average, self-reported sleep time has decreased 1.5–2 h in the USA [43]. Quantitative alterations in sleep duration may impact the metabolic balance of the body, including control of body mass and food intake, glucose metabolism, and adipokine levels (for review, see [44]). In addition to the neurocognitive consequences of sleep loss, recent studies have been focused on the role of sleep in areas outside the brain, including other organs and physiological systems, such as the metabolism [45].

Many studies have shown an association between sleep duration and obesity both in adults and children, suggesting that short sleep duration may be a predictor of weight gain [46–48] and an important risk factor for development of insulin resistance, diabetes, and cardiovascular disease [16,49,50]. A meta-analysis revealed that each reduction of 1 h of sleep per day is associated with an increase of 0.35 kg m^{-2} in body mass index (BMI) [51]. These observed changes due to sleep loss indicate a probable imbalance between food intake and energy expenditure caused by neuroendocrine alterations.

Naturally, sleep is a period of fasting. Glucose utilization by the brain is increased during REM sleep at the end of the night [52], leading to a negative energy balance in the body. However, sleep “resets” the metabolism and energy expenditure rates in the brain, giving effective and flexible control of energy expenditure under changing environmental pressures [53]. Much like sleep, hypothalamic control of metabolism is comprised by mutually inhibiting networks. The appetite-promoting neuropeptide Y (NPY) and agouti-related protein (AGRP) neurons mutually inhibit the appetite-suppressing pro-opiomelanocortin (POMC) and amphetamine-related transcript (CART) neurons. Both sets of neurons work as sensors of the circulating hormones leptin and ghrelin. Leptin is produced by adipose tissues and promotes satiety through inhibition of NPY/AGRP neurons and activation of POMC/CART neurons, with higher levels during sleep compared to awake states, independent of food intake [54]. Recent animal studies have also suggested that leptin participates in sleep regulation, reducing REM sleep and modulating SWS [55]. In turn, ghrelin is an appetite-stimulating hormone produced in the gut, which acts by inhibiting POMC/CART and activating NPY/AGRP. Like leptin, ghrelin has higher levels during sleep, which are followed by a decrease in the morning before the breakfast [56]. Current evidence indicates that ghrelin is also a sleep-promoting factor, able to induce SWS and stimulates GH secretion during the night [57,58].

Sleep curtailment is able to change food intake as a result of decreased secretion of leptin [59–61] and increased secretion of ghrelin [49,59,62], which leads to increased food intake [49]. Two consecutive nights of sleep restriction (4 h of time in bed) in young men were associated with a 28% increase in ghrelin and 18% reduction in leptin during the day, leading to increased hunger (24%) and appetite (23%), mostly for energy-rich foods with high carbohydrate content and low nutritional quality, such as sweets, salty snacks and starchy foods [49]. Six consecutive nights of sleep restriction (4 h of time in bed) increased sympathetic nervous system activity, evening cortisol level and growth hormone, in addition to decreasing glucose effectiveness and the acute insulin response by 30% each, much like is found in non-insulin-dependent diabetes [16]. Buxton and colleagues [63] found that sleep restriction (5 h/night) for 1 week significantly reduced insulin sensitivity, although no correlation was observed with cortisol levels. In a protocol of 14 consecutive days of sleep restriction (5.5 h of time in bed) with *ad libitum* food intake, caloric consumption was increased during the night, when the individual would generally be sleeping, explaining in part the increased vulnerability for weight gain induced by sleep loss [64]. On the other hand, another study did not find differences in hunger ratings after 1 night of total sleep deprivation compared to 1 night of sleep recovery (8 h time in bed) both in men and women [65]. However, actual food intake was not measured in this study, and therefore it is unknown whether participants would or would not have actually increased their food intake during the day of total sleep deprivation compared to the day of sleep rebound [66]. Gonnissen and colleagues [67] evaluated the effect of sleep fragmentation during 8 h on subjective feelings of appetite in men. They did not find a significant reduction in total sleep duration, but rather a reduction in REM sleep and an increase in N2 sleep stage.

The volunteers reported a greater desire to eat after the night of sleep fragmentation, suggesting that sleep quality may be more important than sleep duration for appetite regulation, although they did not measure food intake in this study [67].

Increasing specific clinical evidence has shown that sleep quality and metabolic-related systems are connected. For instance, 50–98% of patients with OSA are morbidly obese [6]. There is an association between OSA and type 2 diabetes [68]. Cross-sectional studies indicate that up to 30% of patients with OSA also present type 2 diabetes, while up to 86% of obese patients with type 2 diabetes have OSA [69,70]. Strong evidence suggests that OSA may increase the risk of developing insulin resistance, glucose intolerance and diabetes [71]. Metabolic disorders and OSA share common pathogenic pathways, such as alterations in autonomic nervous system regulation, increased inflammatory activity, alterations in adipokine levels and endothelial dysfunction, which may be involved in the interplay between these conditions [71]. However, it is not well understood whether these effects are likely due to obesity. In this sense, a systematic review showed that the current literature does not support the hypothesis that OSA independently influences glucose metabolism [72]. The methodological quality varied a lot within the included studies, pointing to a need for more powerful, long-term randomized controlled trials defining changes of insulin resistance as primary endpoint [72].

Obesity is commonly associated with narcolepsy, a sleep disorder characterized by hypocretin (also called orexin) deficiency, excessive daytime sleepiness, and frequent sleep attacks during the day [73]. Narcoleptic patients often present an excess of fat storage in abdominal depots, metabolic alterations, and craving for food with a binge eating pattern [74,75]. The responsiveness of orexin neurons to peripheral metabolic cues, such as leptin and glucose, and the dopaminergic reward system response suggest that both of these 2 neurons are related to the regulation of energy homeostasis and vigilance states [76]. The studies are limited and it is not clear if narcolepsy independently affects glucose metabolism. A recent case-control study showed no clinically relevant pathologic findings in the glucose metabolism of narcoleptic patients compared to weight matched controls [77]. On the other hand, Poli and colleagues showed that narcoleptic patients have higher BMIs and BMI-independent metabolic alterations, such as higher waist circumference, high-density lipoprotein cholesterol, and insulin resistance, compared to idiopathic hypersomnia patients [78]. Evidence has shown that continuous disruption of circadian rhythm in human shift workers is associated with weight gain, metabolic disturbances, type 2 diabetes, and cardiovascular diseases due to increases in postprandial glucose, insulin, cortisol, and mean arterial pressure and decreased 24 h leptin levels [45,79,80].

Another relevant factor that contributes to the development of metabolic disturbances associated with sleep restriction is energy expenditure. Individuals who sleep less are more likely to experience fatigue and sleepiness during the day, which may discourage them from daytime physical activity and promote sedentary behaviors [81,82]. However, the literature presents varied results, in part due to differences in sleep protocols, either total sleep deprivation [83,84]

or partial sleep restriction [85,86], and measurement type (doubly-labeled water [86,87], indirect calorimetry [84], metabolic chamber [83,88], or actigraphy [85,86]). The majority of studies have enrolled small samples with only young, normal-weight men [83,84,88]. Thus, more studies are necessary to determine if sleep duration does affect total energy expenditure and if there is difference between men and women, normal weight and overweight/obese, and young and older individuals. It is possible that improving physical activity could improve sleep, which in turn would impact other components of energy balance, since individual variability in sleep onset latency is reduced by regular physical activity [89].

Not only can sleep affect metabolism, but metabolic changes also can affect sleep architecture [73]. There is an association between late-night-snack intake and sleep patterns [56]. Rodents adjust their sleep onset to match food availability during their sleep-wake cycle [90]. Inversely, food restriction can increase sleep onset latency and reduce total SWS [91]. The common behavior of overeating during a period of sleep deprivation may be a physiological attempt to restore sleep, as it is known that higher food intake promotes sleep [92]. The impact of sleep duration on energy expenditure is less clear due to the multiple factors involved, such as sleeping metabolic rate, thermic effect of food, physical activity, non-exercise activity thermogenesis, etc. [66]. In summary, current literature shows a pattern of increased food intake during periods of sleep loss, mostly in lean and normal sleepers. To date, studies looking for the influence of sleep duration on energy expenditure have produced disparate results due to methodological issues. Due to individual variability, future research assessing whether improving physical activity would improve sleep is also desired for a better understanding.

4. Stress and metabolism

Similarly to sleep, stress is also connected to metabolism. Basal HPA axis activity seems to be dysregulated and overactive both in humans with diabetes and in animal models of type 1 and type 2 diabetes, underlining the neuroendocrine abnormalities common to diabetes-related risk factors such as depression, obesity, hypertension, and cardiovascular diseases [93,94]. Exposure to stressful events leads to increased release of glucocorticoids by activation of the HPA axis [95]. Prolonged activation of the HPA axis may result in maladaptive changes [96], affecting puberty, stature, body composition, as well as leading to obesity, metabolic syndrome, and type 2 diabetes mellitus [97]. Excesses in glucocorticoids increase glucose and insulin and decrease adiponectin levels [98]. Stress exposure alters food intake [99], increasing or decreasing it, depending upon the type of stress [100]. For instance, Ely and colleagues [99] showed that rats subjected to repeated stress by restraint presented increased ingestion of sweet food, while models of chronic variable stress demonstrated a decrease in appetite for sweet food or palatable solutions [101]. There is evidence that glucocorticoids stimulate appetite [102] and increase body weight through the orexigenic effect of NPY [103], an effect

that is inhibited by leptin and insulin [103]. Clinical studies also reveal high food consumption, specifically of palatable food, during periods of psychological stress [104]. The increase in palatable food intake is induced by glucocorticoids [105] and is associated with reward-based eating, as a way to reduce the stress response [106].

Although it appears that hypercortisolemia may contribute to the development of different features of metabolic syndrome, it is not clear in the literature whether glucocorticoids play a role in the pathogenesis of obesity. Some studies show that cortisol levels are not higher in obese subjects, and sometimes they are even lower than in lean subjects [107,108]. This may be, at least in part, a consequence of enhanced cortisol clearance that is thought to accompany obesity, for instance, through increased activity of 5 α -reductase in the liver [108]. Mean 24 h plasmatic ACTH levels were positively correlated with body mass index, reflecting increased hypothalamic drive and reduced negative feedback of cortisol in obesity [109].

Other factors related to cortisol action are also determinants. In this sense, the local expression of 11 β -hydroxysteroid dehydrogenase 1 (11 β -HSD1) plays a role in the relationship between cortisol, adiposity, and metabolic disease [110]. The enzyme 11 β -HSD1, expressed in several peripheral tissues, such as liver and adipose tissue, can modulate HPA axis activity, regenerating active cortisol from its inactive form intracellularly [111]. In humans, 11 β -HSD1 expression is increased in subcutaneous adipose tissue from obese subjects compared to lean subjects [112], being stimulated by TNF α , leptin and adipokines [113,114].

In the presence of insulin, cortisol promotes triglyceride accumulation, mainly in visceral adipocytes, thus leading to increased central adiposity. Masuzaki and colleagues have also demonstrated that overexpression of 11 β -HSD1 in adipose tissue resulted in visceral obesity and metabolic syndrome in mice fed with a high-fat diet [115]. Adipose tissue that overexpressed 11 β -HSD2, the enzyme that inactivates cortisol, protected mice from high-fat diet-induced obesity [116]. The modulation of 11 β -HSD1 might be a promising therapeutic target for obesity and metabolic disturbances. Studies focusing the inhibition of 11 β -HSD1 in animal models of diabetes and obesity have shown improvement of insulin resistance and glucose levels, beyond weight loss [117,118].

Dysregulation of the HPA axis has been associated with some eating disorders [119,120], mainly due to changes in insulin, NPY levels, and other peptides implicated in food intake regulation that can be modulated by cortisol metabolism [112]. Food intake is stimulated by administration of glucocorticoid prednisone in healthy men [121], while diet influences cortisol metabolism, affecting the HPA axis and the reward circuitry for palatable foods [112,122]. Important effects of altered cortisol levels on weight gain are also reported in Cushing's syndrome and Addison's disease, which are both associated with effects such as central obesity/hypercortisolism and weight loss/hypocortisolism, respectively [123].




Fig. 2 – Schematic of the main interactions between sleep, stress and metabolism. Sleep disorders which can lead to sleep loss share common pathways with stress system via HPA axis activation on the metabolic dysfunction, contributing to increased risk of developing obesity and diabetes.

5. Sleep, stress, and metabolism

Because of the new lifestyle imposed by work and family, physical and psychological problems, and social changes due to internet and television, stress and sleep restriction have become endemic, with a major impact on the metabolic process. Importantly, stress hormone levels correlate positively with decreased sleep duration, while both are associated with obesity, metabolic syndrome, and eating disorders [73]. A study by Galvao and colleagues [124] showed that rats subjected to 96 h of paradoxical sleep deprivation present increased immunoreactivity for CRH and orexin as well as higher levels of ACTH and corticosterone, in addition to increased diurnal food intake, but without changes in global food intake. A negative correlation was found between corticosterone and body weight gain throughout paradoxical sleep deprivation [124].

Stress is known to reduce SWS, REM sleep, and delta power, as well as to affect metabolism in rodents, with the magnitude varying according to the type and duration of stress exposure [73]. Sleep deprivation, in turn, activates many stress-related pathways including the HPA axis and sympathetic nervous system, which indirectly modulate arousal and affect the metabolism [26,125]. It has been proposed that the bidirectional relationship between sleep and stress and its impact on metabolism are, in part, mediated by hypocretin circuitry. Hypocretinergic cells project to several CRH-responsive regions in the central nervous system, including *locus coeruleus*, the PVN, the bed nucleus of the *stria terminalis* and the central amygdala [126].

Sleep deprivation *per se* is associated with HPA axis hyperactivity and negatively affects glucose tolerance [16]. The mechanism involved in impaired glucose metabolism following changes in the sleep-wake cycle seems to be the

decreased efficacy of the negative feedback regulation of the HPA axis [42]. Activation of HPA axis may be a risk factor in the development of metabolic syndrome in OSA, via increased visceral obesity, insulin resistance, and sympathetic activity as well as changes in leptin levels [4,127]. However, HPA axis hyperactivity must be only one among several factors that mediate metabolic syndrome in OSA. On the other hand, a recent study in healthy women with clinically diagnosed primary chronic insomnia has demonstrated a dysregulation of circadian cortisol secretion despite normal sleep architecture. Although the limitation of a small number of participants, the authors found that increased midnight cortisol levels were not associated with impaired metabolism of glucose and lipids [128].

The bidirectional interaction between sleep and the HPA axis is complex. Current studies suggest that HPA hyperactivity, sleep loss, and sleep disturbances are closely linked in a vicious circle and play a role in the pathogenesis of metabolic disorders. Understanding sleep and stress system physiology is essential for elucidating the physiopathology of these syndromes and revealing new ways of prevention and treatment.

6. Summary and conclusions

The current review provides evidence for overlap between sleep, stress, and metabolism, which can explain, at least in part, the outcomes observed in the modern society, where sleep deprivation, overnutrition, and chronic exposure to stress potentially lead to the increased incidence and prevalence of metabolic disorders such as obesity and type 2 diabetes. Through hyperactivation of the HPA axis and changes in the neuroendocrine response, sleep loss and chronic stress can lead to metabolic dysfunction. The HPA

axis dysregulation is commonly seen in obesity, sleep deprivation, and sleep disorders such as OSA and insomnia. Conversely, sleep architecture and metabolism are impaired in hypercortisolism conditions such as Cushing's disease, confirming the close relationship between sleep, stress and metabolism, which is summarized in Fig. 2. We conclude that good sleep quality achieved through sleep hygiene and treatment of sleep disorders, in addition to nutritional education with regular meal frequency and circadian alignment of food intake, would be interesting strategies for preventing metabolic disorders. Targeting 11 β -HSD1, a key enzyme in cortisol metabolism in peripheral tissues, and the hypocretin system, which actively and partially regulates the interconnection between sleep, stress and metabolism, might represent a promising therapeutic option for obesity, insulin resistance, and other consequences of excess glucocorticoids which arise from interactions between sleep and stress.

Review criteria

A search for original and review articles that focus on sleep, stress, and metabolism was performed in PubMed. The search terms used were "sleep", "sleep disorders", "sleep loss", "sleep deprivation", "stress", "HPA axis", "cortisol", "corticosterone", "metabolism", "diabetes", "obesity", "glucose", "insulin", "metabolic", "endocrine". We also searched the reference lists of identified articles for further papers. Articles were restricted to human studies. In order to limit the number of references, we selected, whenever possible, a recent review complemented by original papers published after the review.

Acknowledgments

Our studies on this topic have been supported by grants from Associacao Fundo de Incentivo à Pesquisa (AFIP) and Grant #14/15259-2 from São Paulo Research Foundation (FAPESP). M. L.A. and S.T. are recipients of CNPq fellowships (#305177/2013-3 to MLA and #301974/2011-0 to ST). All authors declare no conflicts of interest.

REFERENCES

- [1] Seetho IW, Wilding JP. Sleep-disordered breathing, type 2 diabetes and the metabolic syndrome. *Chron Respir Dis* 2014;11(4):257-75.
- [2] Meerlo P, Sgoifo A, Suchecki D. Restricted and disrupted sleep: effects on autonomic function, neuroendocrine stress systems and stress responsivity. *Sleep Med Rev* 2008;12(3):197-210.
- [3] Spencer RL, Kim PJ, Kalman BA, Cole MA. Evidence for mineralocorticoid receptor facilitation of glucocorticoid receptor-dependent regulation of hypothalamic-pituitary-adrenal axis activity. *Endocrinology* 1998;139(6):2718-26.
- [4] Buckley TM, Schatzberg AF. On the interactions of the hypothalamic-pituitary-adrenal (HPA) axis and sleep: normal HPA axis activity and circadian rhythm, exemplary sleep disorders. *J Clin Endocrinol Metab* 2005;90(5):3106-14.
- [5] Kalsbeek A, van der Spek R, Lei J, Endert E, Buijs RM, Fliers E. Circadian rhythms in the hypothalamo-pituitary-adrenal (HPA) axis. *Mol Cell Endocrinol* 2012;349(1):20-9.
- [6] Spiegel K, Tasali E, Leproult R, Van Cauter E. Effects of poor and short sleep on glucose metabolism and obesity risk. *Nat Rev Endocrinol* 2009;5(5):253-61.
- [7] Steiger A. Neurochemical regulation of sleep. *J Psychiatr Res* 2007;41(7):537-52.
- [8] Gonnissen HK, Hulshof T, Westerterp-Plantenga MS. Chronobiology, endocrinology, and energy- and food-reward homeostasis. *Obes Rev* 2013;14(5):405-16.
- [9] Buxton OM, Copinschi G, Van Onderbergen A, Karrison TG, Van Cauter E. A benzodiazepine hypnotic facilitates adaptation of circadian rhythms and sleep-wake homeostasis to an eight hour delay shift simulating westward jet lag. *Sleep* 2000;23(7):915-27.
- [10] Caufriez A, Moreno-Reyes R, Leproult R, Vertongen F, Van Cauter E, Copinschi G. Immediate effects of an 8-h advance shift of the rest-activity cycle on 24-h profiles of cortisol. *Am J Physiol Endocrinol Metab* 2002;282(5):E1147-53.
- [11] Roth T. Insomnia: definition, prevalence, etiology, and consequences. *J. Clin. Sleep Med* 2007;3(5 Suppl):S7-10.
- [12] Adam K, Tomeny M, Oswald I. Physiological and psychological differences between good and poor sleepers. *J Psychiatr Res* 1986;20(4):301-16.
- [13] Vgontzas AN, Tsigos C, Bixler EO, Stratakis CA, Zachman K, Kales A, et al. Chronic insomnia and activity of the stress system: a preliminary study. *J Psychosom Res* 1998;45(1):21-31.
- [14] Vgontzas AN, Bixler EO, Lin HM, Prolo P, Mastorakos G, Vela-Bueno A, et al. Chronic insomnia is associated with nyctohemeral activation of the hypothalamic-pituitary-adrenal axis: clinical implications. *J Clin Endocrinol Metab* 2001;86(8):3787-94.
- [15] Rodenbeck A, Hajak G. Neuroendocrine dysregulation in primary insomnia. *Rev. Neurol. (Paris)* 2001;157(11 Pt 2):S57-61.
- [16] Spiegel K, Leproult R, Van Cauter E. Impact of sleep debt on metabolic and endocrine function. *Lancet* 1999;354(9188):1435-9.
- [17] Balbo M, Leproult R, Van Cauter E. Impact of sleep and its disturbances on hypothalamo-pituitary-adrenal axis activity. *Int J Endocrinol* 2010;2010:759234.
- [18] Grunstein RR, Stewart DA, Lloyd H, Akinci M, Cheng N, Sullivan CE. Acute withdrawal of nasal CPAP in obstructive sleep apnea does not cause a rise in stress hormones. *Sleep* 1996;19(10):774-82.
- [19] Bratel T, Wennlund A, Carlstrom K. Pituitary reactivity, androgens and catecholamines in obstructive sleep apnoea. Effects of continuous positive airway pressure treatment (CPAP). *Respir Med* 1999;93(1):1-7.
- [20] Tomfohr LM, Edwards KM, Dimsdale JE. Is obstructive sleep apnea associated with cortisol levels? A systematic review of the research evidence. *Sleep Med Rev* 2012;16(3):243-9.
- [21] Rapoport D, Rothenburg SA, Hollander CS, Goldring RM. Obstructive sleep apnea (OSA) alters ultradian rhythm of ACTH secretion. *American Review of Respiratory Disease*. 139 1999:A80.
- [22] Dadoun F, Darmon P, Achard V, Boullu-Ciocca S, Philip-Joet F, Alessi MC, et al. Effect of sleep apnea syndrome on the circadian profile of cortisol in obese men. *Am. J. Physiol. Endocrinol. Metab* 2007;293(2):E466-74.
- [23] Lanfranco F, Gianotti L, Pivetti S, Navone F, Rossetto R, Tassone F, et al. Obese patients with obstructive sleep apnoea syndrome show a peculiar alteration of the corticotroph but not of the thyrotroph and lactotroph function. *Clin Endocrinol (Oxf)* 2004;60(1):41-8.

- [24] Karaca Z, Ismailogullari S, Korkmaz S, Cakir I, Aksu M, Baydemir R, et al. Obstructive sleep apnoea syndrome is associated with relative hypocortisolemia and decreased hypothalamo-pituitary-adrenal axis response to 1 and 250µg ACTH and glucagon stimulation tests. *Sleep Med* 2013;14(2):160-4.
- [25] Weitzman ED, Zimmerman JC, Czeisler CA, Ronda J. Cortisol secretion is inhibited during sleep in normal man. *J Clin Endocrinol Metab* 1983;56(2):352-8.
- [26] Leproult R, Copinschi G, Buxton O, Van Cauter E. Sleep loss results in an elevation of cortisol levels the next evening. *Sleep* 1997;20(10):865-70.
- [27] Follenius M, Brandenberger G, Bandesapt JJ, Libert JP, Ehrhart J. Nocturnal cortisol release in relation to sleep structure. *Sleep* 1992;15(1):21-7.
- [28] Seifritz E, Hemmeter U, Trachsel L, Lauer CJ, Hatzinger M, Emrich HM, et al. Effects of flumazenil on recovery sleep and hormonal secretion after sleep deprivation in male controls. *Psychopharmacology (Berl)* 1995;120(4):449-56.
- [29] Kant GJ, Genser SG, Thorne DR, Pfalser JL, Mougey EH. Effects of 72 hour sleep deprivation on urinary cortisol and indices of metabolism. *Sleep* 1984;7(2):142-6.
- [30] Akerstedt T, Palmblad J, de la Torre B, Marana R, Gillberg M. Adrenocortical and gonadal steroids during sleep deprivation. *Sleep* 1980;3(1):23-30.
- [31] Andersen ML, Bignotto M, Tufik S. Influence of paradoxical sleep deprivation and cocaine on development of spontaneous penile reflexes in rats of different ages. *Brain Res* 2003;968(1):130-8.
- [32] Spath-Schwalbe E, Scholler T, Kern W, Fehm HL, Born J. Nocturnal adrenocorticotropin and cortisol secretion depends on sleep duration and decreases in association with spontaneous awakening in the morning. *J Clin Endocrinol Metab* 1992;75(6):1431-5.
- [33] Hasler G, Buysse DJ, Klaghofer R, Gamma A, Ajdacic V, Eich D, et al. The association between short sleep duration and obesity in young adults: a 13-year prospective study. *Sleep* 2004;27(4):661-6.
- [34] Vorona RD, Winn MP, Babineau TW, Eng BP, Feldman HR, Ware JC. Overweight and obese patients in a primary care population report less sleep than patients with a normal body mass index. *Arch. Intern. Med* 2005;165(1):25-30.
- [35] Ehlers CL, Reed TK, Henriksen SJ. Effects of corticotropin-releasing factor and growth hormone-releasing factor on sleep and activity in rats. *Neuroendocrinology* 1986;42(6):467-74.
- [36] Holsboer F, von Bardeleben U, Steiger A. Effects of intravenous corticotropin-releasing hormone upon sleep-related growth hormone surge and sleep EEG in man. *Neuroendocrinology* 1988;48(1):32-8.
- [37] Gillin JC, Jacobs LS, Fram DH, Snyder F. Acute effect of a glucocorticoid on normal human sleep. *Nature* 1972;237(5355):398-9.
- [38] Born J, DeKloet ER, Wenz H, Kern W, Fehm HL. Gluco- and antiminerocorticoid effects on human sleep: a role of central corticosteroid receptors. *Am. J. Physiol.* 1991;260(2 Pt 1):E183-8.
- [39] Vazquez-Palacios G, Retana-Marquez S, Bonilla-Jaime H, Velazquez-Moctezuma J. Further definition of the effect of corticosterone on the sleep-wake pattern in the male rat. *Pharmacol Biochem Behav* 2001;70(2-3):305-10.
- [40] Born J, Spath-Schwalbe E, Schwachenhofer H, Kern W, Fehm HL. Influences of corticotropin-releasing hormone, adrenocorticotropin, and cortisol on sleep in normal man. *J Clin Endocrinol Metab* 1989;68(5):904-11.
- [41] Bierwolf C, Kern W, Molle M, Born J, Fehm HL. Rhythms of pituitary-adrenal activity during sleep in patients with Cushing's disease. *Exp Clin Endocrinol Diabetes* 2000;108(7):470-9.
- [42] Spiegel K, Knutson K, Leproult R, Tasali E, Van Cauter E. Sleep loss: a novel risk factor for insulin resistance and Type 2 diabetes. *J. Appl. Physiol.* 2005;99(5):2008-19.
- [43] Vioque J, Torres A, Quiles J. Time spent watching television, sleep duration and obesity in adults living in Valencia, Spain. *Int J Obes Relat Metab Disord* 2000;24(12):1683-8.
- [44] Padilha HG, Crispim CA, Zimberg IZ, De-Souza DA, Waterhouse J, Tufik S, et al. A link between sleep loss, glucose metabolism and adipokines. *Braz J Med Biol Res* 2011;44(10):992-9.
- [45] Zimberg IZ, Damaso A, Del Re M, Carneiro AM. H. de Sa Souza, F.S. de Lira, et al., Short sleep duration and obesity: mechanisms and future perspectives. *Cell Biochem Funct* 2012;30(6):524-9.
- [46] Kohatsu ND, Tsai R, Young T, Vangilder R, Burmeister LF, Stromquist AM, et al. Sleep duration and body mass index in a rural population. *Arch Intern Med* 2006;166(16):1701-5.
- [47] Gupta NK, Mueller WH, Chan W, Meininger JC. Is obesity associated with poor sleep quality in adolescents? *Am J Hum Biol* 2002;14(6):762-8.
- [48] Patel SR, Hu FB. Short sleep duration and weight gain: a systematic review. *Obesity (Silver Spring)* 2008;16(3):643-53.
- [49] Spiegel K, Tasali E, Penev P, Van Cauter E. Brief communication: Sleep curtailment in healthy young men is associated with decreased leptin levels, elevated ghrelin levels, and increased hunger and appetite. *Ann Intern Med* 2004;141(11):846-50.
- [50] Taheri S. The link between short sleep duration and obesity: we should recommend more sleep to prevent obesity. *Arch Dis Child* 2006;91(11):881-4.
- [51] Cappuccio FP, Taggart FM, Kandala NB, Currie A, Peile E, Stranges S, et al. Meta-analysis of short sleep duration and obesity in children and adults. *Sleep* 2008;31(5):619-26.
- [52] Maquet P. Sleep function(s) and cerebral metabolism. *Behav Brain Res* 1995;69(1-2):75-83.
- [53] Cirelli C, Faraguna U, Tononi G. Changes in brain gene expression after long-term sleep deprivation. *J Neurochem* 2006;98(5):1632-45.
- [54] Simon C, Gronfier C, Schlienger JL, Brandenberger G. Circadian and ultradian variations of leptin in normal man under continuous enteral nutrition: relationship to sleep and body temperature. *J Clin Endocrinol Metab* 1998;83(6):1893-9.
- [55] Sinton CM, Fitch TE, Gershenfeld HK. The effects of leptin on REM sleep and slow wave delta in rats are reversed by food deprivation. *J Sleep Res* 1999;8(3):197-203.
- [56] Crispim CA, Zalcman I, Dattilo M, Padilha HG, Edwards B, Waterhouse J, et al. The influence of sleep and sleep loss upon food intake and metabolism. *Nutr Res Rev* 2007;20(2):195-212.
- [57] Weikel JC, Wichniak A, Ising M, Brunner H, Friess E, Held K, et al. Ghrelin promotes slow-wave sleep in humans. *Am J Physiol Endocrinol Metab* 2003;284(2):E407-15.
- [58] Schussler P, Uhr M, Ising M, Weikel JC, Schmid DA, Held K, et al. Nocturnal ghrelin, ACTH, GH and cortisol secretion after sleep deprivation in humans. *Psychoneuroendocrinology* 2006;31(8):915-23.
- [59] Spiegel K, Leproult R, L'Hermite-Baleriaux M, Copinschi G, Penev PD, Van Cauter E. Leptin levels are dependent on sleep duration: relationships with sympathovagal balance, carbohydrate regulation, cortisol, and thyrotropin. *J Clin Endocrinol Metab* 2004;89(11):5762-71.
- [60] Taheri S, Lin L, Austin D, Young T, Mignot E. Short sleep duration is associated with reduced leptin, elevated ghrelin, and increased body mass index. *PLoS Med* 2004;1(3):e62.
- [61] Mullington JM, Chan JL, Van Dongen HP, Szuba MP, J. Samaras, N.J. Price, et al., Sleep loss reduces diurnal rhythm

- amplitude of leptin in healthy men. *J Neuroendocrinol* 2003;15(9):851-4.
- [62] Bodosi B, Gardi J, Hajdu I, Szentirmai E, Obal Jr. F, Krueger JM. Rhythms of ghrelin, leptin, and sleep in rats: effects of the normal diurnal cycle, restricted feeding, and sleep deprivation. *Am J Physiol Regul Integr Comp Physiol* 2004;287(5):R1071-9.
- [63] Buxton OM, Pavlova M, Reid EW, Wang W, Simonson DC, Adler GK. Sleep restriction for 1 week reduces insulin sensitivity in healthy men. *Diabetes* 2010;59(9):2126-33.
- [64] Nedeltcheva AV, Kilkus JM. J. Imperial, D.A. Schoeller and P.D. Penev, Insufficient sleep undermines dietary efforts to reduce adiposity. *Ann Intern Med* 2010;153(7):435-41.
- [65] Pejovic S, Vgontzas AN, Basta M, Tsaoussoglou M, Zoumakis E, Vgontzas A, et al. Leptin and hunger levels in young healthy adults after one night of sleep loss. *J Sleep Res* 2010;19(4):552-8.
- [66] St-Onge MP. The role of sleep duration in the regulation of energy balance: effects on energy intakes and expenditure. *J Clin Sleep Med* 2013;9(1):73-80.
- [67] Gonnissen HK, Hursel R, Rutters F, Martens EA, Westerterp-Plantenga MS. Effects of sleep fragmentation on appetite and related hormone concentrations over 24 h in healthy men. *Br J Nutr* 2012;1-9.
- [68] Shaw JE, Punjabi NM, Wilding JP, Alberti KG, Zimmet PZ. E. International Diabetes Federation Taskforce on, et al., Sleep-disordered breathing and type 2 diabetes: a report from the International Diabetes Federation Taskforce on Epidemiology and Prevention. *Diabetes Res Clin Pract* 2008;81(1):2-12.
- [69] Foster GD, Sanders MH, Millman R, Zammit G, Borradaile KE, Newman AB, et al. Obstructive sleep apnea among obese patients with type 2 diabetes. *Diabetes Care* 2009;32(6):1017-9.
- [70] Laaban JP, Daenen S, Leger D, Pascal S, Bayon V, Slama G, et al. Prevalence and predictive factors of sleep apnoea syndrome in type 2 diabetic patients. *Diabetes Metab* 2009;35(5):372-7.
- [71] Lurie A. Metabolic disorders associated with obstructive sleep apnea in adults. *Adv Cardiol* 2011;46:67-138.
- [72] Hecht L, Mohler R, Meyer G. Effects of CPAP-respiration on markers of glucose metabolism in patients with obstructive sleep apnoea syndrome: a systematic review and meta-analysis. *Ger Med Sci*. 9 2011:Doc20.
- [73] Rolls A, Schaich Borg J, and L. de Lecea, Sleep and metabolism: role of hypothalamic neuronal circuitry. *Best Pract Res Clin Endocrinol Metab* 2010;24(5):817-28.
- [74] Kok SW, Overeem S, Visscher TL, Lammers GJ, Seidell JC, Pijl H, et al. Hypocretin deficiency in narcoleptic humans is associated with abdominal obesity. *Obes Res* 2003;11(9):1147-54.
- [75] Fortuyn HA, Swinkels S, Buitelaar J, Renier WO, Furer JW, Rijnders CA, et al. High prevalence of eating disorders in narcolepsy with cataplexy: a case-control study. *Sleep* 2008;31(3):335-41.
- [76] Tsujino N, Sakurai T. Orexin/hypocretin: a neuropeptide at the interface of sleep, energy homeostasis, and reward system. *Pharmacol Rev* 2009;61(2):162-76.
- [77] Beitinger PA, Fulda S, Dalal MA, Wehrle R, Keckeis M, Wetter TC, et al. Glucose tolerance in patients with narcolepsy. *Sleep* 2012;35(2):231-6.
- [78] Poli F, Plazzi G, Di Dalmazi G, Ribichini D, Vicennati V, Pizza F, et al. Body mass index-independent metabolic alterations in narcolepsy with cataplexy. *Sleep* 2009;32(11):1491-7.
- [79] van Drongelen A, Boot CR, Merkus SL, Smid T, van der Beek AJ. The effects of shift work on body weight change - a systematic review of longitudinal studies. *Scand J Work Environ Health* 2011;37(4):263-75.
- [80] Crispim CA, Waterhouse J, Damaso AR, Zimberg IZ, Padilha HG, Oyama LM, et al. Hormonal appetite control is altered by shift work: a preliminary study. *Metabolism* 2011;60(12):1726-35.
- [81] Ortega FB, Chillon P, Ruiz JR, Delgado M, Albers U, Alvarez-Granda JL, et al. Sleep patterns in Spanish adolescents: associations with TV watching and leisure-time physical activity. *Eur J Appl Physiol* 2010;110(3):563-73.
- [82] Knutson KL. Sex differences in the association between sleep and body mass index in adolescents. *J Pediatr* 2005;147(6):830-4.
- [83] Jung CM, Melanson EL, Frydendall EJ, Perreault L, Eckel RH, Wright KP. Energy expenditure during sleep, sleep deprivation and sleep following sleep deprivation in adult humans. *J Physiol*. 2011;589(Pt 1):235-44.
- [84] Benedict C, Hallschmid M, Lassen A, Mahnke C, Schultes B, Schioth HB, et al. Acute sleep deprivation reduces energy expenditure in healthy men. *Am J Clin Nutr* 2011;93(6):1229-36.
- [85] Schmid SM, Hallschmid M, Jauch-Chara K, Wilms B, Benedict C, Lehnert H, et al. Short-term sleep loss decreases physical activity under free-living conditions but does not increase food intake under time-deprived laboratory conditions in healthy men. *Am J Clin Nutr* 2009;90(6):1476-82.
- [86] St-Onge MP, Roberts AL, Chen J, Kelleman M, O'Keefe M, RoyChoudhury A, et al. Short sleep duration increases energy intakes but does not change energy expenditure in normal-weight individuals. *Am J Clin Nutr* 2011;94(2):410-6.
- [87] Nedeltcheva AV, Kilkus JM. J. Imperial, K. Kasza, D.A. Schoeller and P.D. Penev, Sleep curtailment is accompanied by increased intake of calories from snacks. *Am J Clin Nutr* 2009;89(1):126-33.
- [88] Hursel R, Rutters F, Gonnissen HK, Martens EA, Westerterp-Plantenga MS. Effects of sleep fragmentation in healthy men on energy expenditure, substrate oxidation, physical activity, and exhaustion measured over 48 h in a respiratory chamber. *Am J Clin Nutr* 2011;94(3):804-8.
- [89] Buman MP, Hekler EB, Bliwise DL, King AC. Exercise effects on night-to-night fluctuations in self-rated sleep among older adults with sleep complaints. *J. Sleep Res.* 2011;20(1 Pt 1):28-37.
- [90] Fuller PM, Lu J, Saper CB. Differential rescue of light- and food-entrainable circadian rhythms. *Science* 2008;320(5879):1074-7.
- [91] Karklin A, Driver HS, Buffenstein R. Restricted energy intake affects nocturnal body temperature and sleep patterns. *Am J Clin Nutr* 1994;59(2):346-9.
- [92] Penev PD. Sleep deprivation and energy metabolism: to sleep, perchance to eat? *Curr Opin Endocrinol Diabetes Obes* 2007;14(5):374-81.
- [93] Champaneri S, Wand GS, Malhotra SS, Casagrande SS, Golden SH. Biological basis of depression in adults with diabetes. *Curr Diab Rep* 2010;10(6):396-405.
- [94] Beauquis J, Homo-Delarche F, Revsin Y, De Nicola AF, Saravia F. Brain alterations in autoimmune and pharmacological models of diabetes mellitus: focus on hypothalamic-pituitary-adrenocortical axis disturbances. *Neuroimmunomodulation* 2008;15(1):61-7.
- [95] Krolow R, Noschang C, Arcego DM, Huffell AP, Marcolin ML, Benitz AN, et al. Sex-specific effects of isolation stress and consumption of palatable diet during the prepubertal period on metabolic parameters. *Metabolism* 2013;62(9):1268-78.
- [96] Chrousos GP. The role of stress and the hypothalamic-pituitary-adrenal axis in the pathogenesis of the metabolic syndrome: neuro-endocrine and target tissue-related causes. *Int J Obes Relat Metab Disord.* 24 2000;Suppl 2:S50-5.

- [97] Pervanidou P, Chrousos GP. Metabolic consequences of stress during childhood and adolescence. *Metabolism* 2012;61(5):611–9.
- [98] de Oliveira C, de Mattos AB, Biz C, Oyama LM, Ribeiro EB, do Nascimento CM. High-fat diet and glucocorticoid treatment cause hyperglycemia associated with adiponectin receptor alterations. *Lipids Health Dis* 2011;10:11.
- [99] Ely DR, Dapper V, Marasca J, Correa JB, Gamaro GD, Xavier MH, et al. Effect of restraint stress on feeding behavior of rats. *Physiol Behav* 1997;61(3):395–8.
- [100] Varma M, Chai JK, Meguid MM, Gleason JR, Yang ZJ. Effect of operative stress on food intake and feeding pattern in female rats. *Nutrition* 1999;15(5):365–72.
- [101] Willner P. Animal models as simulations of depression. *Trends Pharmacol Sci* 1991;12(4):131–6.
- [102] Dallman MF, Strack AM, Akana SF, Bradbury MJ, Hanson ES, Scribner KA, et al. Feast and famine: critical role of glucocorticoids with insulin in daily energy flow. *Front Neuroendocrinol* 1993;14(4):303–47.
- [103] Stephens TW, Basinski M, Bristow PK, Bue-Valleskey JM, Burgett SG, Craft L, et al. The role of neuropeptide Y in the antiobesity action of the obese gene product. *Nature* 1995;377(6549):530–2.
- [104] Garcia-Prieto MD, Tebar FJ, Nicolas F, Larque E, Zamora S, Garaulet M. Cortisol secretory pattern and glucocorticoid feedback sensitivity in women from a Mediterranean area: relationship with anthropometric characteristics, dietary intake and plasma fatty acid profile. *Clin Endocrinol (Oxf)* 2007;66(2):185–91.
- [105] Pecoraro N, Gomez F, Dallman MF. Glucocorticoids dose-dependently remodel energy stores and amplify incentive relativity effects. *Psychoneuroendocrinology* 2005;30(9):815–25.
- [106] Pecoraro N, Reyes F, Gomez F, Bhargava A, Dallman MF. Chronic stress promotes palatable feeding, which reduces signs of stress: feedforward and feedback effects of chronic stress. *Endocrinology* 2004;145(8):3754–62.
- [107] Rask E, Olsson T, Soderberg S, Andrew R, Livingstone DE, Johnson O, et al. Tissue-specific dysregulation of cortisol metabolism in human obesity. *J Clin Endocrinol Metab* 2001;86(3):1418–21.
- [108] Trivison TG, O'Donnell AB, Araujo AB, Matsumoto AM, McKinlay JB. Cortisol levels and measures of body composition in middle-aged and older men. *Clin Endocrinol (Oxf)* 2007;67(1):71–7.
- [109] Vgontzas AN, Pejovic S, Zoumakis E, Lin HM, Bentley CM, Bixler EO, et al. Hypothalamic-pituitary-adrenal axis activity in obese men with and without sleep apnea: effects of continuous positive airway pressure therapy. *J Clin Endocrinol Metab* 2007;92(11):4199–207.
- [110] Anagnostis P, Athyros VG, Tziomalos K, Karagiannis A, Mikhailidis DP. Clinical review: The pathogenetic role of cortisol in the metabolic syndrome: a hypothesis. *J Clin Endocrinol Metab* 2009;94(8):2692–701.
- [111] Tomlinson JW, Walker EA, Bujalska IJ, Draper N, Lavery GG, Cooper MS, et al. 11beta-hydroxysteroid dehydrogenase type 1: a tissue-specific regulator of glucocorticoid response. *Endocr Rev* 2004;25(5):831–66.
- [112] Bose M, Olivan B, Laferrere B. Stress and obesity: the role of the hypothalamic-pituitary-adrenal axis in metabolic disease. *Curr Opin Endocrinol Diabetes* 2009;16(5):340–6.
- [113] Tomlinson JW, Moore J, Cooper MS, Bujalska I, Shahmanesh M, Burt C, et al. Regulation of expression of 11beta-hydroxysteroid dehydrogenase type 1 in adipose tissue: tissue-specific induction by cytokines. *Endocrinology* 2001;142(5):1982–9.
- [114] Dieudonne MN, Sammari A, Dos Santos E, Leneuve MC, Giudicelli Y, Pecquery R. Sex steroids and leptin regulate 11beta-hydroxysteroid dehydrogenase I and P450 aromatase expressions in human preadipocytes: Sex specificities. *J Steroid Biochem Mol Biol* 2006;99(4-5):189–96.
- [115] Masuzaki H, Paterson J, Shinyama H, Morton NM, Mullins JJ, Seckl JR, et al. A transgenic model of visceral obesity and the metabolic syndrome. *Science* 2001;294(5549):2166–70.
- [116] Kershaw EE, Morton NM, Dhillon H, Ramage L, Seckl JR, Flier JS. Adipocyte-specific glucocorticoid inactivation protects against diet-induced obesity. *Diabetes* 2005;54(4):1023–31.
- [117] Alberts P, Engblom L, Edling N, Forsgren M, Klingstrom G, Larsson C, et al. Selective inhibition of 11beta-hydroxysteroid dehydrogenase type 1 decreases blood glucose concentrations in hyperglycaemic mice. *Diabetologia* 2002;45(11):1528–32.
- [118] Sundbom M, Kaiser C, Bjorkstrand E, Castro VM, Larsson C, Selen G, et al. Inhibition of 11betaHSD1 with the S-phenylethylaminothiazolone BVT116429 increases adiponectin concentrations and improves glucose homeostasis in diabetic KKAY mice. *BMC Pharmacol.* 8 2008:3.
- [119] Gluck ME. Stress response and binge eating disorder. *Appetite* 2006;46(1):26–30.
- [120] Laue L, Gold PW, Richmond A, Chrousos GP. The hypothalamic-pituitary-adrenal axis in anorexia nervosa and bulimia nervosa: pathophysiological implications. *Adv Pediatr* 1991;38:287–316.
- [121] Tataranni PA, Larson DE, Snitker S, Young JB, Flatt JP, Ravussin E. Effects of glucocorticoids on energy metabolism and food intake in humans. *Am. J. Physiol.* 1996;271(2 Pt 1):E317–25.
- [122] Stimson RH, Johnstone AM, Homer NZ, Wake DJ, Morton NM, Andrew R, et al. Dietary macronutrient content alters cortisol metabolism independently of body weight changes in obese men. *J Clin Endocrinol Metab* 2007;92(11):4480–4.
- [123] Lucassen EA, Cizza G. The Hypothalamic-Pituitary-Adrenal Axis, Obesity, and Chronic Stress Exposure: Sleep and the HPA Axis in Obesity. *Curr Obes Rep* 2012;1(4):208–15.
- [124] Galvao Mde O, Sinigaglia-Coimbra R, Kawakami SE, Tufik S, Suchecki D. Paradoxical sleep deprivation activates hypothalamic nuclei that regulate food intake and stress response. *Psychoneuroendocrinology* 2009;34(8):1176–83.
- [125] Meerlo P, Koehl M, van der Borgh K, Turek FW. Sleep restriction alters the hypothalamic-pituitary-adrenal response to stress. *J Neuroendocrinol* 2002;14(5):397–402.
- [126] Winsky-Sommerer R, Yamanaka A, Diano S, Borok E, Roberts AJ, Sakurai T, et al. Interaction between the corticotropin-releasing factor system and hypocretins (orexins): a novel circuit mediating stress response. *J Neurosci* 2004;24(50):11439–48.
- [127] Sandoval DA, Davis SN. Leptin: metabolic control and regulation. *J Diabetes Complications* 2003;17(2):108–13.
- [128] Seelig E, Keller U, Klarhofer M, Scheffler K, Brand S, Holsboer-Trachsler E, et al. Neuroendocrine regulation and metabolism of glucose and lipids in primary chronic insomnia: a prospective case-control study. *PLoS One* 2013;8(4):e61780.